

Phase diffusion of a two-component Bose-Einstein condensates: exact and short-time solutions for arbitrary coherent spin state

G. R. Jin, B. B. Wang, and Y. W. Lu

Department of Physics, Beijing Jiaotong University, Beijing 100044, China

We investigate phase diffusion of a two-component Bose-Einstein condensates prepared initially in arbitrary coherent spin state $|\theta_0, \phi_0\rangle$. Analytical expression of the phase-diffusion time is presented for $\theta_0 \neq \pi/2$ case. In comparison with the symmetrical case (i.e., $\theta_0 = \pi/2$), we find that the diffusion process becomes slowly due to the reduced atom number variance.

PACS numbers: 03.75.Mn, 05.30.Jp, 42.50.Lc

Phase diffusion of Bose-Einstein condensates (BECs) [1–7], aroused from atom-atom interactions destroys phase coherence, and thus sets a limit to the applications of the condensates in high-precision measurement and quantum information processing. The phase diffusion is observable by measuring fringe visibility in atomic interference experiments [8–12]. For a two-component BECs prepared initially in a coherent spin state (CSS) $|\theta_0, \phi_0\rangle$ with $\theta_0 = \pi/2$ [see Ref. [13], or Eq. (2)], it has been shown that the single-particle coherence (or fringe visibility) decays exponentially with the time scale $\chi t_d = (N/2)^{-1/2}$ [2–5], dependent upon atom number N and the self-interaction strength $\chi \sim (a_{11} + a_{22} - 2a_{12})/2$. Here, a_{ii} and a_{ij} ($i \neq j$) denote intra- and inter-species s-wave scattering lengths, respectively. In this paper, we generalize previous works by considering arbitrary initial CSS, i.e., $\theta_0 \neq \pi/2$. Exact and short-time solutions of the single-particle coherence are obtained. In comparison with $\theta_0 = \pi/2$ case, we find that the diffusion becomes slowly due to relatively narrow atom number variance.

Considering a two-component BECs with internal states $|1\rangle$ and $|2\rangle$ confined in a deep potential, we adopt single-mode approximation (SMA) [14], i.e., keeping the condensed-mode wave function $\Phi_0(r)$ for the two components, so the total system can be well described by second-quantized Hamiltonian ($\hbar = 1$):

$$\hat{H} = \omega_1 \hat{n}_1 + \omega_2 \hat{n}_2 + u_{12} \hat{n}_1 \hat{n}_2 + \frac{u_{11}}{2} (\hat{b}_1^\dagger)^2 (\hat{b}_1)^2 + \frac{u_{22}}{2} (\hat{b}_2^\dagger)^2 (\hat{b}_2)^2, \quad (1)$$

where $\hat{b}_i$, $\hat{b}_i^\dagger$, and $\hat{n}_i$ ($= \hat{b}_i^\dagger \hat{b}_i$) are the annihilation, creation, and number operators for the internal states $i = 1, 2$, respectively. In addition, ω_i are single-particle energies, and $u_{ij} = (4\pi a_{ij}/M) \int d^3r |\Phi_0(r)|^4$ are atom-atom interaction strengthes. Particle number operator $\hat{N} = \hat{n}_1 + \hat{n}_2$ is a conserved quantity and is set to the c number N . Hamiltonian (1) can be rewritten as $\hat{H} = \delta \hat{J}_z + \chi \hat{J}_z^2$, where the detuning $\delta = \omega_2 - \omega_1 + (u_{22} - u_{11})(N-1)/2$, and the self-interaction strength $\chi = (u_{11} + u_{22} - 2u_{12})/2$. The spin operators $\hat{J}_+ = (\hat{J}_-)^\dagger = \hat{b}_2^\dagger \hat{b}_1$ and $\hat{J}_z = (\hat{n}_2 - \hat{n}_1)/2$, obeying SU(2) algebra. The nonlinearity $\chi \hat{J}_z^2$ has been proposed to prepare spin squeezed state [15–17] and also quantum entangled state [18, 19]. Note that the model

considered here with nonzero δ can be also used to study the BECs in an asymmetric double-well [20]. Squeezing via coupling of the BECs in a double-well potential with a cavity light field has been investigated in Ref. [21].

Besides the spin squeezing, mean-field interaction $\chi \hat{J}_z^2$ also leads to phase diffusion in the two-component BECs [1–7, 12], which can be illuminated schematically by Husimi Q function $Q(\theta, \phi; t) = |\langle \theta, \phi | \Psi(t) \rangle|^2$, where

$$|\theta, \phi\rangle = \exp\{i\theta(\hat{J}_x \sin \phi - \hat{J}_y \cos \phi)\} |j, j\rangle \quad (2)$$

is arbitrary CSS [13] and $|\Psi(t)\rangle = e^{-i\hat{H}t} |\Psi(0)\rangle$ is a state vector at any time t . For an initial state $|\Psi(0)\rangle = |\theta_0, \phi_0\rangle$, analytic expression of the Q function reads [13]

$$Q(\theta, \phi; 0) = |\langle \theta, \phi | \theta_0, \phi_0 \rangle|^2 = \left[\frac{1 + \cos \Theta}{2} \right]^{2j}, \quad (3)$$

where $\cos \Theta = \cos \theta \cos \theta_0 + \sin \theta \sin \theta_0 \cos(\phi - \phi_0)$. The Q function can be plotted in three-dimensional phase space (i.e., Bloch sphere) [15, 16], or alternatively, in a two-dimensional phase space (ϕ, s_z) via a mapping $s_z = j \cos \theta$ [12, 17, 22]. As shown in Fig. 1(a) and Fig. 1(b), the density of $Q(\theta, \phi; 0)$ is distributed isotropically, indicating the minimal uncertainty relationship of the initial CSS. Under the government of nonlinear interaction $\chi \hat{J}_z^2$, the system will evolve into a spin squeezed state [15] with anisotropic distribution of the Q function [see Fig. 1(c)]. As time increases, the spin system is over-squeezed [15, 23] and shows a spread of the Q function along the ϕ axis, which simulates an increased relative phase fluctuation (i.e., phase diffusion) [12], as shown in Fig. 1(d).

It was shown that the most sensitive states to the diffusion are the symmetrical CSS with $\theta_0 = \pi/2$, corresponding to equal populations between the two internal states [7]. The phase-diffusion time scale is $\chi t_d = 1/\sqrt{j}$ (with $j = N/2$) [2–5]. Here, we generalize it for arbitrary initial CSS

$$|\Psi(0)\rangle = |\theta_0, \phi_0\rangle = \sum_{m=-j}^j c_m |j, m\rangle, \quad (4)$$




FIG. 1: Husimi Q function $Q(\theta, \phi; t)$ for initial CSS $|\theta_0 = \pi/2, \phi_0 = 0\rangle$ on Bloch sphere (a), with Bloch vector $\vec{s} = \langle \vec{J} \rangle = j(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$ and $j = N/2 = 30$. The Q functions in the phase space (ϕ, s_z) for various times: (b) $t = 0$, (c) $t = t_s = 0.0764\chi^{-1}$, and (d) $t = 2t_d = 0.2651\chi^{-1}$. Time scale $\chi t_s \simeq 3^{1/6}(2j)^{-2/3}$ is for optimal spin squeezing [15–17], and $\chi t_d = j^{-1/2}$ is phase-diffusion time [2–5]. The Q function is normalized by its maximum value.

where the amplitudes

$$c_m = \binom{2j}{j-m}^{1/2} \cos^{j+m}(\frac{\theta_0}{2}) \sin^{j-m}(\frac{\theta_0}{2}) e^{i(j-m)\phi_0} \quad (5)$$

In single-particle picture, the CSS corresponds to all the atoms occupying a superposed state [15]: $\cos(\theta_0/2)|2\rangle + e^{i\phi_0} \sin(\theta_0/2)|1\rangle$, where the polar angle θ_0 and the azimuth angle ϕ_0 determine atom population and the relative phase between the two internal states, respectively. In Fig. 2, we plot $|c_m|^2$ as a function of m . One can find that the probability distribution of the CSS can be treated as a Gaussian wave packet [see below Eq. (11)]. At any time t , the spin system evolves into

$$|\Psi(t)\rangle = \sum_{m=-j}^j c_m e^{-i(\delta m + \chi m^2)t} |j, m\rangle, \quad (6)$$

i.e., a superposition of atomic number state $|j, m\rangle = |j-m\rangle_1 |j+m\rangle_2$. Due to the presence of atom-atom interaction ($\chi \neq 0$), each number state has different phase evolution rate, which in turn lead to collapse and revival of the Rabi oscillation, a phenomenon that is well-known in quantum optics [14]. Similar effect has been investigated in two-component BECs [24], atom-molecular BECs [25–27], and exciton emission [28–30].

Phase diffusion of the BECs considered here is in fact collapse of the first-order temporal correlation function [3, 7]:

$$g_{12}^{(1)} = \frac{|\rho_{12}^{(1)}|}{\sqrt{\rho_{11}^{(1)} \rho_{22}^{(1)}}} \equiv \frac{|\langle \hat{J}_+ \rangle|}{\sqrt{j^2 - \langle \hat{J}_z \rangle^2}}, \quad (7)$$

where $\rho_{kl}^{(1)} = \langle \hat{b}_k^\dagger \hat{b}_l \rangle / N$ with $k, l = 1, 2$ are the elements of the single-particle density matrix [7]. The coherence $g_{12}^{(1)}$

is observable in experiments by extracting the visibility of the interference fringes [8–12]. The expectation value $\langle \hat{J}_z \rangle = \langle \Psi(t) | \hat{J}_z | \Psi(t) \rangle$, reads

$$\langle \hat{J}_z \rangle = \sum_{m=-j}^j m |c_m|^2 = j \cos \theta_0, \quad (8)$$

which is a constant for a fixed polar angle of the initial CSS θ_0 . This is because of conserved operator $\hat{J}_z$ with respect to the Hamiltonian $\hat{H}$. Similarly, we obtain the expectation value $\langle \hat{J}_z^2 \rangle = j^2 \cos^2 \theta_0 + (j/2) \sin^2 \theta_0$, and

$$\begin{aligned} \langle \hat{J}_+ \rangle &= e^{i(\phi_0 + \delta t + \chi t)} \cot \frac{\theta_0}{2} \sum_{m=-j}^j (j-m) |c_m|^2 e^{2im\chi t} \\ &= j e^{i(\phi_0 + \delta t)} \sin \theta_0 [\cos \chi t + i \cos \theta_0 \sin \chi t]^{2j-1} \quad (9) \end{aligned}$$

where we have used the relation: $c_{m+1} = (j-m)^{1/2}(j+m+1)^{-1/2} \cot(\theta_0/2) e^{-i\phi_0} c_m$. Inserting Eq. (8) and Eq. (9) into Eq. (7), we further obtain the exact solution of the coherence

$$g_{12}^{(1)}(t) = [1 - \sin^2(\theta_0) \sin^2(\chi t)]^{j-1/2}, \quad (10)$$

which shows a decay of the coherence, i.e., phase diffusion. Such a dephasing process depends sensitively on the self-interaction strength χ and the polar angle θ_0 of the initial CSS; while the detuning δ and the azimuth angle ϕ_0 gives *vanishing* contribution to the coherence.




FIG. 2: Probability distribution, $|c_m|^2$ as a function of m for the initial CSS $|\theta_0, \phi_0\rangle$. (a) $\theta_0 = \pi/2$ and (b) $\theta_0 = \pi/3$. The empty circle is given by Eq. (5), and solid line is given by Eq. (11). The parameter $j = N/2 = 30$.

Time scale of the diffusion is determined by short-time behavior of the coherence. Following Imamoglu et al. [3], we treat the initial CSS as a Gaussian wave-packet with its peak located at $m \simeq \langle \hat{J}_z \rangle = j \cos \theta_0$, namely

$$|c_m|^2 \simeq \frac{1}{[2\pi(\Delta \hat{J}_z)^2]^{1/2}} \exp \left[-\frac{(m - \langle \hat{J}_z \rangle)^2}{2(\Delta \hat{J}_z)^2} \right], \quad (11)$$

where the width

$$(\Delta \hat{J}_z)^2 = \langle \hat{J}_z^2 \rangle - \langle \hat{J}_z \rangle^2 = (j/2) \sin^2 \theta_0. \quad (12)$$

Due to conserved particle number $N (= 2j)$, $\hat{n}_1 = j - \hat{J}_z$ and $\hat{n}_2 = j + \hat{J}_z$, and thus atom number variances $(\Delta \hat{n}_1)^2 = (\Delta \hat{n}_2)^2 = (\Delta \hat{J}_z)^2$ [17].

In Fig. 2, we check the validity of Eq. (11) by comparing it with the exact result of $|c_m|^2$ [where c_m is given by Eq. (5)], and find both results fit with each other. For symmetric BEC with $\theta_0 = \pi/2$, we have $\langle \hat{J}_z \rangle = 0$ and atom number variance $(\Delta \hat{J}_z)^2 = j/2$; while for $\theta_0 \neq \pi/2$, the variance (width) becomes narrow, as shown in Fig. 2(b). Inserting Eq. (11) into Eq. (9), and replacing the discrete sum over m by an integral, we get

$$\begin{aligned} \langle \hat{J}_+ \rangle &\simeq \frac{1}{\sqrt{\pi}} e^{i(\phi_0 + \delta t + \chi t)} e^{2i\langle \hat{J}_z \rangle \chi t} \cot \frac{\theta_0}{2} \\ &\times \int_{\rho_{\min}}^{\rho_{\max}} [j - \langle \hat{J}_z \rangle - \sqrt{2}(\Delta \hat{J}_z)\rho] \\ &\times \exp[-\rho^2 + 2i\sqrt{2}\chi t(\Delta \hat{J}_z)\rho] d\rho, \end{aligned} \quad (13)$$

where we have set

$$\rho = \frac{m - \langle \hat{J}_z \rangle}{\sqrt{2}(\Delta \hat{J}_z)}. \quad (14)$$

For $\theta_0 \sim \pi/2$ and $j \rightarrow \infty$ (large N limit), the integral upper limit $\rho_{\max} = \sqrt{j} \tan(\theta_0/2) \rightarrow \infty$ and the lower limit $\rho_{\min} = -\sqrt{j} \cot(\theta_0/2) \rightarrow -\infty$. In the short-time limit ($\chi t \ll 1$), $1 - 2i\chi t \cos^2(\theta_0/2) \simeq \exp\{-2i\chi t \cos^2(\theta_0/2)\}$, and thus from Eq. (13) we obtain

$$\begin{aligned} \langle \hat{J}_+ \rangle &\simeq j \sin(\theta_0) \exp\{i[\phi_0 + \delta t + \chi t(2j - 1) \cos \theta_0]\} \\ &\times \exp\left[-2\chi^2(\Delta \hat{J}_z)^2 t^2\right], \end{aligned} \quad (15)$$

where two integrals: $\int_{-\infty}^{\infty} e^{-\rho^2} e^{i\lambda\rho} d\rho = \sqrt{\pi} e^{-\lambda^2/4}$ and $\int_{-\infty}^{\infty} \rho e^{-\rho^2} e^{i\lambda\rho} d\rho = i\lambda\sqrt{\pi} e^{-\lambda^2/4}/2$ have been used in derive of the above result. Thus, the short-time solution of the coherence reads

$$g_{12}^{(1)}(t) \simeq \exp\left[-2\chi^2(\Delta \hat{J}_z)^2 t^2\right] \equiv \exp\left[-(t/t_d)^2\right], \quad (16)$$

with characteristic time scale of the phase diffusion

$$\chi t_d = \frac{1}{\sqrt{2}(\Delta \hat{J}_z)} = \frac{1}{\sqrt{j} \sin \theta_0}. \quad (17)$$

Obviously, the phase-diffusion time $\chi t_d = 1/\sqrt{j}$ for the initial CSS with $\theta_0 = \pi/2$ [3, 7, 12]; while for $\theta_0 \neq \pi/2$, our results show that the diffusion becomes slowly due to an enhanced factor $\sin \theta_0$ in the phase-diffusion rate [see also Fig. 3]. The revival of the coherence occurring later at a time $\chi t_r = \pi$ [5, 7] is observable in real experiment, such as $t_r = \pi/\chi \simeq 108.7$ microsecond for nonlinearity $\chi \simeq 2\pi \times 4.6$ Hz and particle number $N = 60$ [12].

Finally, we note that similar results of Eq. (9) and Eq. (15) have been obtained in Ref. [31]. However, the authors focused on the increase of the diffusion time as $\chi \rightarrow 0$, but not $\theta_0 \neq \pi/2$. In addition, we emphasize that for negative χ the so-called phase separation may take place due to dynamically unstable of the BECs [32–39]. Both the phase diffusion and the phase separation




FIG. 3: Time evolution of the first-order temporal coherence $g_{12}^{(1)}(t)$ for various polar angles: $\theta_0 = \pi/6$ (dotted line, squares), $\pi/4$ (dashed line, triangles), and $\pi/2$ (solid line, circles). The empty squares, triangles, and circles are given by Eq. (10) for corresponding θ_0 's, while the curves are given by Eq. (16). Horizontal grid line gives the value $1/e$, vertical lines denote the diffusion times t_d for different θ 's. Other parameters are $N = 60$, $\chi = 2\pi \times 4.6$ Hz [12].

reduce the Ramsey fringes' visibility (i.e., the first-order coherence $g_{12}^{(1)}$). However, the latter effect results from dynamics of spatial degree-of-freedom, rather than that of the internal states considered here.

In summary, we have investigated phase diffusion of two-component BEC for arbitrary initial CSS. We show analytically that for the CSS with $\theta_0 \neq \pi/2$, the diffusion process is suppressed due to the reduced atom number variance $(\Delta \hat{J}_z)^2$ below the standard quantum limit $j/2$ [see also Eqs. (12) and (17)]. Our analytic results are based upon the SMA, which works well to describe the condensates tightly confined in a three-dimensional (3D) trapping potential. For a lower dimensional cases [40], the coherence $g_{12}^{(1)}$ [or the fringe visibility] is expected to decay more quickly due to the whole continuum of excitation modes in confined degrees of freedom [12, 41].

Acknowledgments

This work is supported by the NSFC (Contract No. 10804007).

-
- [1] Lewenstein M and You L 1996 *Phys. Rev. Lett.* **77** 3489
 - [2] Wright E M, Walls D F, and Garrison J C 1996 *Phys. Rev. Lett.* **77** 2158
 - [3] Imamoglu A, Lewenstein M, and You L 1997 *Phys. Rev. Lett.* **78** 2511.
 - [4] Javanainen J and Wilkens M 1997 *Phys. Rev. Lett.* **78** 4675; Javanainen J and Ivanov M. Y. 1999 *Phys. Rev. A* **60** 2351
 - [5] Castin Y and Dalibard J 1997 *Phys. Rev. A* **55** 4330
 - [6] Law C K, Pu H, Bigelow N P, and Eberly J H 1998 *Phys. Rev. A* **58** 531

- [7] Khodorkovsky Y, Kurizki G, and Vardi A 2008 *Phys. Rev. Lett.* **100** 220403
- [8] Orzel C, Tuchman A K, Fenselau M L, Yasuda M, and Kasevich M A 2001 *Science* **291** 2386; Greiner M, Mandel O, Esslinger T, Hänsch T W, Bloch I 2002 *Nature* **415** 39; Li W, Tuchman A K, Chien H-C, and Kasevich M A 2007 *Phys. Rev. Lett.* **98** 040402
- [9] Schumm T, Hofferberth S, Andersson L M, Wildermuth S, Groth S, Bar-Joseph I, Schmiedmayer J, Krüger P 2005 *Nature Phys.* **1** 57; Shin Y, Sanner C, Jo G-B, Pasquini T A, Saba M, Ketterle W, Pritchard D E, Vengalattore M, and Prentiss M 2005 *Phys. Rev. A* **72** 021604(R)
- [10] Chuu C-S, Schreck F, Meyrath T P, Hanssen J L, Price G N, and Raizen M G 2005 *Phys. Rev. Lett.* **95** 260403
- [11] Jo G-B, Shin Y, Will S, Pasquini T A, Saba M, Ketterle W, Pritchard D E, Vengalattore M, and Prentiss M 2007 *Phys. Rev. Lett.* **98** 030407
- [12] Widera A, Trotzky S, Cheinet P, Fölling S, Gerbier F, Bloch I, Gritsev V, Lukin M D, and Demler E 2008 *Phys. Rev. Lett.* **100** 140401
- [13] Radcliffe J M 1971 *J. Phys. A* **4** 313; Arecchi F T, Courtens E, Gilmore R, and Thomas H 1972 *Phys. Rev. A* **6** 2211
- [14] Walls D F and Milburn G J 2008 *Quantum Optics* (the 2nd edition, Springer-Verlag, Berlin Heidelberg), and references therein
- [15] Kitagawa M and Ueda M 1993 *Phys. Rev. A* **47** 5138; Sørensen A, Duan L M, Cirac J I, Zoller P 2001 *Nature* **409** 63
- [16] Jin G R and Kim S W 2007 *Phys. Rev. A* **76** 043621; *ibid.* 2007 *Phys. Rev. Lett.* **99** 170405
- [17] Jin G R and Law C K 2008 *Phys. Rev. A* **78** 063620
- [18] Wei W 2005 *Chinese Physics A* **14** 2407
- [19] Cheng R and Liang J-Q 2007 *Chinese Physics A* **16** 834
- [20] Wang D-L, Wang F-J, Xiao Y-F and Yan X-H 2006, *Acta Physica Sinica* **55** 547
- [21] Kong L-B, Zhan M-S, and Zhou L 2008, *Chinese Physics B* **17** 1601
- [22] Trimborn F, Witthaut D, and Korsch H J 2009 *Phys. Rev. A* **79** 013608
- [23] Shalm L K, Adamson R B A, and Steinberg A M 2009 *Nature* **457** 67
- [24] Kuang L M and Ouyang Z W 2000 *Phys. Rev. A* **61** 023604; Kuang L M and Zhou L 2003 *Phys. Rev. A* **68** 043606; Li W-D, Zhou X-J, Wang Y-Q, Liang J-Q, and Liu W-M 2001 *Phys. Rev. A* **64** 015602
- [25] Javanainen J and Mackie M 1999, *Phys. Rev. A* **59**, R3186; Vardi A, Yurovsky V A, and Anglin J R 2001, *Phys. Rev. A* **64** 063611
- [26] Jin G R, Kim C K, and Nahm K 2005 *Phys. Rev. A* **72** 045602
- [27] Cai W, Jing H, Zhan M-S, Xu J-J 2007 *Chin. Phys. Lett.* **24** 867
- [28] Jin G R and Liu W M 2004 *Phys. Rev. A* **70** 013803; Jin G R, Liang Z X, and Liu W M 2004 *J. Opt. B: Quantum Semiclass. Opt.* **6** 296
- [29] Glazov M M and Kavokin K V 2006 *Phys. Rev. B* **73** 245317
- [30] Li W, Song P, Liu J, Yang X 2007 *J. Phys. B* **40** 1957
- [31] Sinatra A and Castin Y 2000 *Eur. Phys. J. D* **8** 319
- [32] Ao P and Chui S T 1998 *Phys. Rev. A* **58** 4836; Wu Biao and Niu Qian 2001, *Phys. Rev. A* **64** 061603(R); Kono-top V V and Salerno M 2002 *Phys. Rev. A* **65** 021602(R)
- [33] Kasamatsu K and Tsubota M 2004 *Phys. Rev. Lett.* **93** 100402; Kasamatsu K and Tsubota M 2006 *Phys. Rev. A* **74** 013617
- [34] Li L, Li Z, Malomed B A, Mihalache D, Liu W M 2005 *Phys. Rev. A* **72** 033611; Jin G R, Kim C K, and Nahm K 2005 *Phys. Rev. A* **72** 045601
- [35] Belmonte-Beitia J, Pérez-García V M, Vekslerchik V 2007 *Chaos, Solitons and Fractals*, **32** 1268
- [36] Huang Jin-Song, Chen Hai-Feng, and Xie Zhen-Wei 2008 *Acta Phys. Sinica* **57** 3435
- [37] Sorokina E A, Maimistova A I and Elyutin S O 2009 *Physica D* **238** 1394
- [38] Alon O E, Streltsov A I, and Cederbaum L S 2009 *Phys. Rev. A* **79** 022503
- [39] Baizakov B B, Bouketir A, Messikh A, and Umarov B A 2009 *Phys. Rev. E* **79** 046605
- [40] Cui H-T, Wang L-C, and Yi X-X 2004 *Acta Physica Sinica* **53** 991
- [41] Bistritzer R and Altman E 2007 *PNAS* **104** 9955; Burkov A A, Lukin M D, and Demler E 2007 *Phys. Rev. Lett.* **98** 200404.